


Double-blind, Placebo-controlled, randomized, multicenter study evaluating Tafasitamab in combination with Lenalidomide and Rituximab versus Placebo plus Lenalidomide and Rituximab



Primary end point

- PFS in FL population

Secondary end points

- PFS and MRD-negativity rate in FL and MZL population
- OS, PET-CR, best ORR, DoR, QoL in overall and FL population
- Safety

Select inclusion criteria

- Age ≥18
- Histologically confirmed Grade 1-3a FL, or nodal, splenic or extra nodal MZL
- ECOG PS 0-2
- Previously treated with at least 1 systemic anti-CD20 immunotherapy or chemo-immunotherapy
- Documented relapsed, refractory or PD after treatment with systemic therapy
- Ability and willingness to receive prophylaxis/treatment of thromboembolic events

Select exclusion criteria

- Histology other than FL or MZL, or clinical evidence of transformed lymphoma
- Prior non-hematologic malignancy
- CNS lymphoma involvement
- Congestive heart failure
- Active systemic infection
- Prior use of Lenalidomide plus Rituximab
- Systemic anti-lymphoma/investigational therapy within 28 days prior cycle 1

The efficacy and safety of the investigational compound discussed^a have not been established. There is no guarantee that this compound^a will become commercially available for the use(s) under investigation.

For more information, visit [IncyteClinicalTrials.com](https://www.incyte.com/clinicaltrials), contact us at +800 00027423 or by email at clintrials@incyte.com, or click on the QR code on the right to visit the trial on clinicaltrials.gov

For reporting adverse events, contact eumedinfo@incyte.com or +49 8007239013

